

$\mathbb{R}$ $\mathbb{U}$ M E^x
 E H_ψ $\mathbb{C}$ $M \neq \psi$ ψ $=$ E

'E[˘] ĭ HΨ η r Ġ 'E ρ à‰ ~ 'E T
 X à κ Â

[illegible]
$$\begin{aligned} & \Psi \quad \mathcal{H}\Psi \mathcal{H} \quad \quad \quad \tilde{\omega} \quad \mathcal{H} \quad \hat{A} \quad \mathcal{L}\Psi \quad \mathcal{H}\gamma \quad \quad \quad = \\ & \quad \gamma \, \tilde{\eta} = \quad ' \\ & \bar{\omega} \, \tilde{\alpha} \Psi \quad \gamma \quad \quad \quad \tilde{\omega} \quad \quad \quad = \quad \hat{e}^{\tau} \, \mathcal{G} \\ & \quad \tilde{E}^{\tilde{\alpha}} \quad \quad \quad \mathcal{L} \quad \mathcal{H} \quad \uparrow \quad \quad \quad \tilde{\omega} \, \hat{A} \, \hat{e}^{\tau} \, \mathcal{G} \quad \quad \quad \tilde{E}^{\tilde{\alpha}} \quad \quad \quad \mathcal{L} \quad \mathcal{H} \\ & \uparrow \quad \quad \quad \tilde{E} \quad \tilde{\omega} \, \eta \quad \quad \quad \mathcal{G} \\ & \mathcal{H} \, \tilde{\alpha} \Psi \quad \gamma \quad \quad \quad \tilde{\omega} \quad \quad \quad = \quad \hat{e}^{\tau} \, \mathcal{G} \, \mathcal{A} \\ & \mathcal{L} \quad \quad \quad \tilde{\omega} \quad \sim \mathcal{L} \quad \quad \quad \tilde{\omega} \, \hat{A} \, \hat{e}^{\tau} \, \mathcal{G} \, \mathcal{A} \quad \mathcal{L} \\ & \quad \tilde{\omega} \quad \sim \mathcal{L} \quad \quad \quad \tilde{E} \quad \tilde{\omega} \, \eta \quad \quad \quad \mathcal{G} \end{aligned}$$
 $\hat{E} \hat{A}$

γ_{12} $\tilde{0}$ M $E^{\tilde{\alpha}}$ $H\psi$
 $H\dot{n}H$ $\ddot{X}$ $H\gamma$